



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

POINT SETS AND CREMONA GROUPS. PART III

By Arthur A. Coble

DEPARTMENT OF MATHEMATICS, JOHNS HOPKINS UNIVERSITY

Received by the Academy, August 23, 1916

In Part I¹ of this series projectively distinct sets P_n^k of n points in S_k were mapped upon points of a space $\Sigma_{k(n-k-2)}$ and a certain Cremona group G_n in Σ was obtained by permutation of the points of the set. In Part II² the G_n appeared as merely a subgroup of a more important group $G_{n,k}$ in $\Sigma_{k(n-k-2)}$ which also is defined by P_n^k . In particular the G_6 in Σ_4 attached to P_6^2 is a subgroup of the $G_{6,2}$ in Σ_4 which has the order 51840 and is isomorphic with the group of the lines on a cubic surface.

The purpose of this Part III is to utilize the $G_{6,2}$ in the problem of determining the lines of a cubic surface C^3 . It appears that there is a one-to-one correspondence between the invariants of C^3 and the invariant spreads of $G_{6,2}$ in Σ_4 . The lines of C^3 can be rationally expressed in terms of a solution of the form problem of $G_{6,2}$ by means of a typical representation of C^3 in the hexahedral form with the aid of the linear covariants of C^3 . In order to solve the form problem of $G_{6,2}$ the simplest linear system of irrational invariants of C^3 is employed. This system is of dimension 9 and the members appear in Σ_4 as quintic spreads. Under the invariant subgroup $\Gamma_{6,2}$ of $G_{6,2}$ of index two this linear system separates into two skew linear systems each of dimension 4 with the important property that the members of the two systems are permuted under the operations of $G_{6,2}$ precisely as the points and S_3 's of a linear space S_4 are permuted under the elements of a correlation group in S_4 whose collineation subgroup is the Burkhardt group G_{25920} in S_4 . The form problem of $G_{6,2}$ can then be solved in terms of a solution of the form problem of G_{25920} by using the point invariants of G_{25920} and in addition five invariants of G_{25920} linear in the S_3 coördinates and of degrees 1, 7, 9, 13, 15 in the point coördinates.

The method for solving the form problem of G_{25920} is suggested by the properties of the normal hyperelliptic surface M_2^{18} of grade 3 in S_8 obtained parametrically by using 9 linearly independent theta functions of the third order and zero characteristic. The M_2^{18} admits a collineation group $G_{2,81}$ which contains 81 involutions. If I is one of these involutions the fixed S_3 and fixed S_4 of I meet M_2^{18} in 6 and 10 points respectively. The M_2^{18} is projected from the fixed S_4 upon the fixed S_3 into a doubly covered Weddle surface and from the fixed S_3 upon the fixed S_4 into a doubly covered 2-way N_2^6 which has a node

at α . If N_2^6 be projected from α it becomes a Kummer surface. There is a family of ∞^3 surfaces M_2^{18} with the same $G_{2,81}$. By projection we obtain a family of $N_2^{6'}$'s whose node α runs over a quartic spread J_4 —the simplest invariant of G_{25920} . The 10 points in the S_4 of I run over the Hessian J_{10} of J_4 . The spread J_4 is its own Steinerian and the polar cubic of a point α on J_4 as to J_4 is a Segre cubic spread with nodes at the 10 points on J_{10} , and of course a simple point at α . The point α on its polar cubic determines a binary sextic—the fundamental sextic of the hyperelliptic functions. In this way the solution of the form problem of G_{25920} in terms of hyperelliptic modular functions becomes apparent at once *in the special case* when $J_4 = 0$. This restriction is removed later by a conventional method. The conclusions above all are drawn from the existence of a set of 9 quadrics whose complete intersection is the normal spread M_2^{18} and whose coefficients are the modular forms α .

The above determination of the lines of C^3 differs from that of Klein¹ in that no equation of degree 27 or other resolvent equation is employed. All the processes are effected within the domain of the invariants and linear covariants of C^3 . Klein also uses as fundamental form problem that of the Maschke collineation group in S_3 rather than the Burkhardt form problem. This implies the isolation of a root of the underlying binary sextic. The accessory irrationalities required are thereby somewhat simpler.

¹ These PROCEEDINGS, 1, 245 (1915); *Trans. Amer. Math. Soc.*, 16, 155 (1915). This series of investigations has been pursued under the auspices of the Carnegie Institution of Washington, D. C.

² These PROCEEDINGS, 2, 244 (1916); *Trans. Amer. Math. Soc.*, 17, 345 (1916).

³ That an equation of degree 27 for the lines of a cubic surface could be solved by hyperelliptic modular functions was first pointed out by Klein, *J. Math., Paris*, Ser. 4, 4, 169 (1888). His suggestions were elaborated by Witting, *Math. Ann., Leipzig*, 29, 167 (1887); by Maschke, *Ibid.*, 33, 317 (1889); and by Burkhardt, *Ibid.*, 35, 198 (1890), 38, 161 (1891), 41, 313 (1893).

THE INTERFERENCES OF SPECTRA BOTH REVERSED AND INVERTED

By Carl Barus

DEPARTMENT OF PHYSICS, BROWN UNIVERSITY

Received by the Academy, September 6, 1916

This is an interesting combination of the two methods of investigation hitherto given (Carnegie Publications, No. 249, 1916, §4) and not very difficult to produce. Retaining the adjustment for inverted spectra